

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Blood loss in orthotopic liver transplantation: a retrospective analysis of transfusion requirements and the effects of autotransfusion of cell saver blood in 164 consecutive patients	Hendriks HG, van der Meer J, Klomp maker IJ, Choudhury N, Hagenaa rs JA, Porte RJ, de Kam PJ, Slooff MJ, de Wolf JT	Department of Anaesthesiology, Groningen University Hospital, The Netherlands	Blood Coagul Fibrinolysis 2000 Apr;11 Suppl 1:S87-93
Determination of eicosanoid and cytokine production in salvaged blood, stored red blood cell concentrates, and whole blood	Jacobi KE, Wanke C, Jacobi A, Weisbach V, Hemmerling TM	Department of Anesthesiology, University of Erlangen-Nuremberg, Erlangen, Germany	J Clin Anesth 2000 Mar; 12(2):94-9
Potassium as a surrogate marker of debris in cell-salvaged blood	Szpisjak DF, Edgell DS, Bissonnette B	Departments of Anesthesia and Perfusion, The Hospital for Sick Children, University of Toronto, Ontario, Canada	Anesth Analg 2000 Jul;91(1):40-3
Laparoscopic surgery with intraoperative autologous blood transfusion in patients with heavy hemoperitoneum due to ectopic pregnancy [In Process Citation]	Yamada T, Kasamatsu H	Department of Pathology, Osaka Medical College, 2-7 Daigaku-machi, Takatsuki, Osaka 569-8686, Japan	J Am Assoc Gynecol Laparosc 2000 May;7(2):255-6
Autoapheresis and intraoperative blood salvage in oncologic surgery	Valbonesi M, Bruni R, Lercari G, Florio G, Carlier P, Morelli F	Immunohematology Services, San Martino University Hospital, Genova, Italy	Transfus Sci 1999 Oct;21(2):129-39
The effect of intraoperative autotransfusion on antibiotic pharmacokinetics during elective abdominal aortic aneurysm repair	Aiono S, Stacey A, Magee TR, Galland RB	Department of Surgery, Royal Berkshire Hospital, Reading, UK	Cardiovasc Surg 2000 Mar;8(2):137-40
Plasma levels of eicosanoids after transfusion of intraoperatively salvaged blood	Jacobi K, Walther A, Lorler H, Neidhardt B	Klinik fur Anesthesiologie, Chirurgische Klinik mit Poliklinik, Friedrich-Alexander-Universitat Erlangen-Nurnberg, Erlangen Deutschland	Vox Sang 2000;78(1):31-6
The results of intraoperative autotransfusion in orthopaedic surgery	Benli IT, Akalin S, Duman E, Citak M, Kis M	1 st Department of Orthopaedics and Traumatology, Ankara Social Security Hospital, Turkey	Bull Hosp Jt Dis 1999;58(4); 184-7
Autotransfusion performed on a patient with cis AB blood group	Kawahito S, Kitahata H, Kimura H, Tanaka K, Oshita S	Department of Anaesthesiology, Tokushima University School of Medicine, Japan	Br J Anaesth 1999 Sep;83(3): 491-2
Cell salvage in the Jehovah's Witness patient [letter; comment]	Waters JH, Potter PS		Anesth Analg 2000 Jan;90(1): 229-30
Primary and revision total hip replacement in patients who are Jehovah's Witnesses	Nelson CL, Stewart JG	University of Arkansas for Medical Sciences, Little Rock 72205, USA	Clin Orthop 1999 Dec;(360): 251-61
The impact of intraoperative autologous blood transfusion during type III radical hysterectomy for early-stage cervical cancer	Mirhashemi R, Averette HE, Deepika K, Estape R, Angioli R, Martin J, Rodriquez M, Penalver MA	Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, University of Miami School of Medicine/Jackson Memorial Medical Center 33136, USA	Am J Obstet Gynecol 1999 Dec; 181(6):1310-5; discussion 1315-6
Minimizing venous air embolism from reinfusion bags [letter; comment]	Benumof JL		Anesthesiology 1999 Dec;91(6): 1962

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
The Ross operation in a Jehovah's Witness: a paradigm for heart surgery in children without transfusion	Miyaji K, Hannan RL, Ojito JW, White JA, Burke RP	Division of Cardiovascular Surgery, Miami Children's Hospital, Florida 33155-4069, USA	Ann Thorac Surg 2000 Mar; 69(3):935-7
Impact of red cell salvage on transfusion requirements during elective abdominal aortic aneurysm repair	Szalay D, Wong D, Lindsay T	Division of Vascular Surgery, The Toronto Hospital and The University of Toronto, Toronto, Ontario, Canada	Ann Vasc Surg 1999 Nov;13(6): 576-81
Use of autologous transfusion in the management of acute traumatic haemothorax in the accident and emergency department	McGhee A, Swinton S, Watt M	Accident and Emergency Department, Monklands District General Hospital, Lanarkshire	J Accid Emerg Med 1999 Nov; 16(6):451-2
Infected and malignant fields are an absolute contraindication to intraoperative cell salvage: fact or fiction?	Thomas MJ	Blood Care Foundation, UK	Transfus Med 1999 Sep;9(3): 269-78
Autotransfusion in coronary artery bypass grafting; disparity in laboratory tests and clinical performance	Flom-Halvorsen HI, Ovrum E, Tangen G, Brosstad F, Ringdal MA, Oystese R	Department of Cardiac Surgery, Oslo Heart Center, University of Oslo, Rikshospitalet, Oslo, Norway	J Thorac Cardiovasc Surg 1999 Oct;118(4)610-7
Troponin I and autotransfusion of shed mediastinal blood in redo cardiac operation [letter; comment]	Chen LW		Ann Thorac Surg 1999 Aug; 68(2):629-30
Effects of unprocessed and processed cardiopulmonary bypass blood retransfused into patients after cardiac surgery	Walpoth BH, Eggensperger N, Hauser SP, Neidhart P, Kurt G, Spaeth PJ, Althaus U	Department of Cardiovascular Surgery, Anesthesiology, Central Hematology Laboratory, University Hospital, Insel, Berne, Switzerland	Int J Artif Organs 1999;22(4): 210-6
Autotransfusion of washed shed mediastinal fluid decreases the requirement for autologous blood transfusion following cardiac surgery: a prospective randomized trial	Dalrymple-Hay MJ, Pack L, Deakin CD, Shephard S, Ohri SK, Haw MP, Livesay SA, Monro JL	Wessex Cardiothoracic Centre, Department of Cardiothoracic Surgery, Southampton General Hospital, UK	Eur J Cardiothorac Surg 1999 Jun;15(6):830-4
Influence of different autotransfusion devices on the quality of salvaged blood	Reents W, Babin-Ebell J, Misoph MR, Schwarzkopf A, Elert O	Department of Cardiothoracic Surgery, University Hospital, Wurzburg, Germany	Ann Thorac Surg 1999 Jul; 68(1):58-62
Amniotic fluid removal during cell salvage in the cesarean section patient [see comments]	Waters JH; Biscotti C, Potter PS, Phillipson E	Departments of General Anesthesiology, Obstetrics and Gynecology, and Anatomic Pathology, Cleveland Clinic Foundation, Cleveland, Ohio 44195, USA	Anesthesiology 2000 Jun;92(6): 1531-6
Efficacy of postoperative blood salvage following total hip arthroplasty in patients with and without deposited autologous units [In Process Citation]	Grosvenor D, Goyal V, Goodman S	Division of Orthopaedic Surgery, Stanford University Medical Center, California 94305, USA	J Bone Joint Surg Am 2000 Jul; 82-A(7):951-4 [MEDLINE record in process]
Comparison of blood-conservation strategies in cardiac surgery patients at high risk for bleeding	Nuttall GA, Oliver WC, Ereth MH, Santrach PJ, Bryant SC, Orszulak TA, Schaff HV	Department of Anesthesiology, Mayo Clinic, Rochester, Minnesota 55905, USA	Anesthesiology 2000 Mar;92(3): 674-82
Inflammatory cytokines and their receptors in arterial and mixed venous blood before, during and after infusion of drained untreated blood	Krohn CD, Reikeras O, Aasen AO	National Hospital, Centre of Orthopaedics, Trondhjemsveien 132, 0570 Oslo, Norway	Transfus Med 1999 June;9(2): 125-30

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Perioperative blood transfusions: indications and options	McFarland JG	Blood Center of Southeastern Wisconsin, Milwaukee 53201-2178, USA	Chest 1999 May;115(5 Suppl): 113S-121S
Safety and efficacy of autologous platelet transfusion in cardiac surgery: comparison of cryopreservation, blood collection on the day before surgery, and blood collection during surgery	Yokomuro M, Ebine K, Shiroma K, Tamura S, Kumabe S, Ohtuki M, Suzuki H, Uchida S	Department of Cardiovascular Surgery, Toho University School of Medicine, Ohashi Hospital, Ohashi 2-17-6, Meguro-ku, Tokyo, 153-8515, Japan	Cyrobiology 1999 May;38(3): 236-42
Complement activation and increased systemic and pulmonary vascular resistance indices during infusion of postoperatively drained untreated blood	Krohn CD, Reikeras O, Mollnes TE	National Hospital, Centre for Orthopaedics, Oslo, Norway	Br J Anaesth 1999 Jan;82(1): 47-51
Quality of reinfused drainage blood after total knee arthroplasty	Jensen CM, Pilegaard R, Hviid K, Nielsen JD, Nielsen HJ	Department of Orthopaedic Surgery, Copenhagen Country Hospital, University of Copenhagen, Gentofte, Denmark	J Arthroplasty 1999 Apr;14(3): 312-8
Intraoperative autotransfusion in small children; an in vitro investigation to study its feasibility	Booke M, Hagemann O, Van Aken H, Erren M, Wullenweber J, Bone HG	Department of Anesthesiology and Intensive Care, University of Munster, Germany	Anesth Analg 1999 Apr;88(4): 763-5
Formation of complement split products and proinflammatory cytokines by reinfusion of shed autologous blood	Bengtsson A, Avall A, Hyliner M, Bengtson JP	Department of Anaesthesiology, Oregon Health Sciences University, Portland, USA	Toxicol Lett 1998 Nov 23; 100-101:129-33
Blood separation with two different autotransfusion devices: effects on blood cell quality and coagulation variables	Rosolski T, Matthey T, Frick U, Hachenberg T	Department of Anesthesiology and Intensive Care Medicine, Ernst-Moritz-Arndt – University, Greifswald – Germany	Int J Artif Organs 1998 Dec; 21(12):820-4
Application of cell-salvage during cesarean section	Potter PS, Waters JH, Burger GA, Mraovic B	Department of General Anesthesiology, Cleveland Clinic Foundation, Ohio 44195, USA	Anesthesiology 1999 Feb;90(2): 619-21
Autotransfusion supported by erythropoietin therapy in transurethral resection of the prostate	Goya N, Toda F, Nishino S, Tokumoto T, Iguchi Y, Kobayashi H, Ohshima T, Kihara T, Tanabe K, Yagisawa T, Nakazawa H, Toma H, Fujii H	Department of Urology, Tokyo Women's Medical University, Japan	Scand J Urol Nephrol 1998 Dec; 32(6):383-7
Audit of autotransfusion in spine surgery	Shulman G, Solanki DR, Nicodemus CL, Flores IM, Hadjipavlou AB	Department of Pathology, University of Texas Medical Branch, Galveston 77555-0192, USA	Int Orthop 1998;22(5):303-7
Massive bleeding during spine surgery in a patient with ankylosing spondylitis	Tetzlaff JE, Yoon HJ, Bell G	Department of General Anesthesiology, Cleveland Clinic Foundation, Ohio 44195, USA	Can J Anaesth 1998 Sep;45(9): 903-6

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
The safety of intraoperative autologous blood collection and autotransfusion during cesarean section	Rebarber A, Lonser R, Jackson S, Copel JA, Sipes S	Department of Obstetrics and Gynecology, Yale University School of Medicine, New Haven, Connecticut, USA	Am J Obstet Gynecol 1998 Sep; 179(3 Pt 1):715-20
Methyl methacrylate levels in unwashed salvage blood following unilateral total knee arthroplasty	Hand GC, Henderson M, Mace P, Sherif N, Newman JH, Goldie DJ	Avon Orthopaedic Centre, Southmead Hospital, Bristol, United Kingdom	J Arthroplasty 1998 Aug;13(5): 576-9
Postoperative blood salvage in hip and knee arthro-plasty. A prospective study on cost effectiveness in 161 patients	Rizzi L, Bertacchi P, Ghezzi LM, Bellavita P, Scudeller G	Department of Orthopedics, Matteo Rota Orthopedic Institute, Bergamo, Italy	Acta Orthop Scand 1998 Feb; 69(1):31-4
Blood salvage during caesarean section [see comments]	Rainaldi MP, Tazzari PL, Scagliarini G, Borghi B, Conte R	Anaesthesia and Intensive Care Maternity Unit, Policlinico S. Orsola, Bologna, Italy	Br J Anaesth 1998 Feb;80(2): 195-8
Are you ready for bloodless medicine?	Wolfe S		RN2000 May;63(5):42-6
Autologous Platelet Gel with Growth Factors in Facial Plastic Surgery	Adler SC		August 2000
Using Platelet-Rich Plasma to Develop an Autologous Membrane for Growth Factor Delivery in Dental Implant Therapy	Garg AK		Dental Implantology, Volume 11, Number 6, June 2000
Quantification of growth factor levels using a simplified method of platelet-rich plasma gel preparation	Landesberg R, Roy M, Glickman RS	New York University/Bellevue Hospital Center, NY, USA	J Oral Maxillofac Surg 2000 Mar; 58(3):297-300; discussion 300-1
Quality of harvested autologous platelets compared with stored donor platelets for use after cardiopulmonary bypass procedures	Crowther M, Ford I, Jeffrey RR, Urbaniak SJ, Greaves M	Department of Medicine and Therapeutics, University of Aberdeen, Aberdeen and Northeast of Scotland Blood Transfusion Service, and Cardiothoracic Surgery, Aberdeen	Br J Haematol 2000 Oct; 111(1): 175-181
Incidence and risk factors for allogenic blood transfusion during major joint replacement using an integrated autotransfusion regimen. The Rizzoli Study Group on Orthopaedic Anaesthesia	Borghi B, Casati A	Department of Anaesthesiology, IRCCS Rizzoli Orthopaedic Institute, Bologna and Department of Anaesthesiology, IRCCS H San Raffaele, Via Olgettina 60-20132 Milan, Italy	Eur J Anaesthesiol 2000; 17(7): 411-7
Reinfusion of mediastinal blood after heart surgery	Martin J, Robitaille D, Perrault LP, Pellerin M, Page P, Searle N, Cartier R, Hebert Y, Pelletier LC, Thaler HT, Carrier M	Departments of Surgery and Anesthesia and the Laboratory of Hematology, Montreal Heart Institute and University of Montreal, Montreal, Quebec, Canada	J Thorac Cardiovasc Surg 2000 Sep;120(3):499-504
Intraoperative autotransfusion for repair of unruptured aneurysms of the infrarenal abdominal aorta. A multicenter study of 203 patients. Association Universitaire de Recherche en Chirurgie (AURC)	Patra P, Chaillou P, Bizouarn P	Service de Chirurgie Vasculaire, Hopital Guillaume et Rene Laennec, Nantes, France	J Cardiovasc Surg (Torino) 2000 Jun;41(3):407-13

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Effect of autotransfusion on fibrinolysis in open heart patients	Warsaw DS, Niewiarowska A, Theman T	Department of Surgery, St. Luke's Hospital, Bethel, PA 18015, USA	J Extra Corpor Technol 2000 Mar;32(1):20-4
Tissue factor antigen and activity in serum of postoperatively shed blood used for autologous transfusion [In Process Citation]	Krohn CD, Reikeras O, Bjornsen S, Brosstad F	National Hospital, Centre for Orthopaedics, Oslo, Norway	Blood Coagul Fibrinolysis 2000 Apr;11(3):219-23
Use of interventions to minimize perioperative allogeneic blood transfusion in Australia. A survey by the International Study of Perioperative Transfusion (ISPOT) Study Group	Henry DA, Henderson KM, Fryer JL, Treloar CJ, McGrath KM, Deveridge SF	Faculty of Medicine and Health Services, University of Newcastle	Med J Aust 2000 Apr 17;172(8):365-9
Platelet-rich plasma sequestration, with therapeutic platelet yields, reduces allogeneic transfusion in complex cardiac surgery	Stover EP, Siegel LC, Hood PA, O'Riordan GE; McKenna TR	Department of Anesthesia, Stanford University School of Medicine, Stanford, California 94305, USA	Anesth Analg 2000 mar;90(3): 509-16
Transfusion of washed and centrifuged shed RBCs during maxillofacial surgery affects cytokine concentrations	Rohling RG, Zimmerman AP, Zellweger R, Schanz U	Departments of Cranio-Maxillofacial Surgery, Trauma Surgery, and Internal Medicine (Division of Hematology), Institute of Anesthesiology, University Hospital, Zurich, Switzerland	Transfusion 2000 40:1352-6
Retransfusion of suctioned blood during cardiopulmonary bypass impairs hemostatis	DeHaan J, Boonstra PW, Monnik SH, Ebels T, van Oeveren W	Department of Cardiothoracic Surgery, University Hospital, Groningen, The Netherlands	Ann Thorac Surg 1995 Apr; 59(4):901-7
Processing scavenged blood with a cell saver reduces cerebral lipid microembolization	Kincaid EH, Jones TJ, Stump DA, Brown WR, Moody DM, Deal DD, Hammon JW	Department of Cardiothoracic Surgery, Wake Forest University School of Medicine, Winston-Salem, North Carolina 27157, USA	Ann Thorac Surg 2000 Oct; 70(4):1296-300
The ability of the Haemonetics 4 Cell Saver System to remove tissue factor from blood contaminated with amniotic fluid	Bernstein HH, Rosenblatt MA, Gettes M, Lockwood C	Department of Anesthesiology, Mount Sinai Medical Center, New York, NY 10029-6574, USA	Anesth Analg 1997 Oct;85(4): 831-3
Cell Salvage and Leukodepletion	Cross MH	Department of Anesthesia, The General Infirmary at Leeds, Leeds, UK	Perfusion 2001; Volume 16 (suppl):61-66
Cell saver, ultrafiltration and direct transfusion: comparative study of three blood processing techniques	Eichert I, Isgro F, Kiessling AH, Saggau W	Department of Cardiac Surgery, Ludwigshafen, Germany	Thorac Cardiovasc Surg 2001 Jun;49(3):149-52
Debris elimination from partially-filled cell salvage bowls	Szpisjak DF	Department of Anesthesiology, The National Naval Medical Center, 8901 Wisconsin Ave., Bethesda, MD 20889, USA	Anesth Analg 2001 May;92(5):1137-8
Does cell-saver blood administration and free hemoglobin load cause renal dysfunction?	Klodell CT, Richardson JD, Bergamini TM, Spain DA	Department of Surgery, University of Louisville School of Medicine and the Veterans Administration Medical Center, Kentucky 40292, USA	Am Surg 2001 Jan;67(1):44-7

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Postoperative blood salvage and reinfusion in spinal surgery: blood quality, effectiveness and impact on patient blood parameters	Sebastian C, Romero R, Olalla E, Ferrer C, Garcia-Vallejo JJ, Munoz M	Department of Orthopaedic Surgery, Parque San Antonio Hospital, Malaga, Spain	Eur Spine J 2000 Dec;9(6): 458-65
Smart suction device for less blood trauma: a comparison with Cell Saver	Mueller XM, Tevaearai HT, Horisberger J, Augstburger M, Boone Y, von Segesser LK	Clinic for Cardiovascular Surgery and Surgical Intensive Care Unit, Centre Hospitalier Universitaire Vaudois, CH-1011 Lausanne, Switzerland	Eur J Cardiothorac Surg 2001 Apr;19(4):507-11
Processing scavenged blood with a cell saver reduces cerebral lipid microembolization	Kincaid EH, Jones TJ, Stump DA, Brown WR, Moody DM, Deal DD, Hammon JW Jr	Department of Cardiothoracic Surgery, Wake Forest University School of Medicine, Winston-Salem, North Carolina 27157, USA	Ann Thorac Surg 2000 Oct; 70(4):1296-300
Intra-operative blood salvage in abdominal trauma: a review of 5 years' experience	Hughes LG, Thomas DW, Wareham K, Jones JE, John A, Rees M	Intensive Care Unit, Swansea NHS Trust, Morriston Hospital, Swansea SA6 6NL, UK	Anaesthesia 2001 Mar;56(3): 217-20
Effect of Partial Filling Autotransfusion Bowls on the Quality of the Reinfused Product	Tremain KD, Stammers AH, et al	Division of Clinical Perfusion Education, School of Allied Health Professions, University of Nebraska Medical Center, Omaha, NE	J Extra Corpor Technol 2001 May;33(2):80-5
Platelet-rich plasma application in sinus graft surgery: Part I—Background and processing techniques	Lozada JL, Caplanis N, Proussaefs P, Willardsen J, Kammeyer G	Department of Restorative Dentistry, School of Dentistry, Loma Linda University, Loma Linda, CA 92350, USA	J Oral Implantol 2001;27(1): 38-42
Complement split products and pro-inflammatory cytokines in salvaged blood after hip and knee arthroplasty	Andersson I, Tylman M, Bengston JP, Bengston A	Department of Anesthesiology & Intensive Care, East Hospital/Sahlgrenska University Hospital, Gothenburg, Sweden	Can J Anaesth 2001 Mar;48(3): 251-5
Fat elimination from autologous blood	Booke M, Van Aken H, Storm M, Fritsche F, Wirtz S, Hinder F	Klinik und Poliklinik fur Anesthesiologie und operative Intensivmedizin, University of Munster, 48129, Munster, Germany	Anesth Analg 2001 Feb;92(2): 341-3
Unwashed shed blood infusion causes deterioration in right ventricular function after coronary artery surgery	Kitano T, Hattori S, Miyakawa H, Yoshitake S, Iwasaka H, Noguchi T	Department of Anesthesiology, Oita Medical University, Oita-gun, Oita, Japan	Anaesth Intensive Care 2000 Dec;28(6):642-5
A Cytokine Analysis of the Effect of Cell Saver on Blood in Coronary Bypass Surgery	Sandoval S, Airawi S, Samee M, Satheesan R	Lutheran & Maimondes Medical Center Research Institute; Department of Surgery, Division of Cardiothoracic Surgery; Lutheran Surgical Department, Brooklyn, NY, USA	The Heart Surgery Forum #2001-19264/4(2):113-9

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Different preparation methods to obtain platelet components as a source of growth factors for local application	Zimmermann R, Jakubietz R, Jakubietz M, Strasser E, Schlegel A, Wiltfang J, Eckstein R	Departments of Transfusion Medicine and Hemostaseology and of Oral and Maxillofacial Surgery, Friedrich Alexander University, Erlangen-Nurnberg, Erlangen, Germany	Transfusion 2001 Oct;41(10): 1217-24
Does Perioperative Autologous-blood-component Therapy Demonstrate Clinical Benefits in Patients Undergoing Orthopedic Surgery	Jutzi H, Herren H, Rehle R, Steiner J, Ueltschi P		The Internet Journal of Anesthesiology 1998 Vol2N2
Intraoperative cell salvage in radical retropubic prostatectomy(1)	Gray CL, Amling CL, Polston GR, Powell CR, Kane CJ	Department of Urology, Naval Medical Center, San Diego, California, USA	Urology 2001 Nov;58(5):740-5
Autotransfusion in cardiac surgery	Cross MH	Department of Anaesthesia, The General Infirmary at Leeds, UK	Perfusion 2001 Sep;16(5): 391-400
A cytokine analysis of the effect of cell saver on blood in coronary bypass surgery	Sandoval S, Alrawi S, Samee M, Satheesan R, Raju R, Cunningham J, Acinapura A	Lutheran & Maimonides Medical Center Research Institute, Brooklyn, NY, USA	Heart Surg Forum 2001;4(2): 113-7; discussion 117-9
The biology of platelet-rich plasma	Schmitz J, Hollinger J		J Oral Maxillofac Surg 2001 Sep;59(9):1119-21
Are leukocytes in salvaged washed autologous blood harmful for the recipient? The results of a pilot study.	Innerhofer P, Wiedermann FJ, Tiefenthaler W, Schobersberger W, Klingler A, Velik-Salchner C, Oswald E, Salner E, Irschick E	Department of Anesthesia and Critical Care Medicine, The Leopold-Franzens University of Innsbruck, Innsbruck, Austria	Anesth Analg 2001 Sep;93(3): 566-72
Quantitative microbiologic study of blood salvaged by intraoperative membrane filtration	Nosanchuk, J	Department of Pathology and Laboratory Medicine, Cayuga Medical Center at Ithaca, Ithaca, NY 14850, USA	Arch Pathol Lab Med 2001 Sep; 125(9):1204-6
[Bacterial contamination of salvaged blood in open heart surgery: is that an airborne contamination or a normal skin flora contamination?]	Ishida T, Nakano S, Nakatani H, Gomi A, Sato T, Saegusa N, Ito A, Okada A, Tazawa Y	Department of Cardiovascular Surgery, Kanto Medical Center NTT EC, Tokyo, Japan	Kyobu Geka 2001 Aug;54(9): 753-7 [Article in Japanese]
Release of interleukin-10 by reinfusion of salvaged blood after knee arthroplasty	Tylman M, Bengtson J, Avall A, Hyliner M, Bengtsson A	Department of Anesthesiology and Intensive Care, Sahlgrenska University Hospital, Ostra, 41685 Gothenburg, Sweden	Intensive Care Med 2001 Aug; 27(8):1379-84
Effect of abciximab on the outcome of emergency coronary artery bypass grafting after failed percutaneous coronary intervention	Singh M, Nuttall G, Ballman K, Mullany C, Berger P, Holmes D Jr, Bell M	Division of Cardiovascular Diseases and Internal Medicine, Mayo Clinic, Rochester, MN 55905, USA	Mayo Clin Proc 2001 Aug; 76(8):784-8
Emergency coronary artery bypass grafting: does excessive preoperative anticoagulation increase bleeding complications and transfusion requirements?	Grubitzsch H, Wollert H, Eckel L	Heart and Diabetes Centre Mecklenburg-Vorpommern, Clinic of Cardiothoracic and Vascular Surgery, Karlsburg, Germany	Cardiovasc Surg 2001 Oct; 9(5):510-6

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Increased interleukin-6 in collected drainage blood after total knee arthroplasty: an association with febrile reactions during retransfusion	Handel M, Winkler J, Hornlein R, Northoff H, Heeg P, Teschner M, Sell S	Section of Orthopaedic Surgery, University Hospital, Tübingen, Germany	Acta Orthop Scand 2001 Jun; 72(3):270-2
Effects of autologous fresh platelet concentrate on haemostasis in cardiac reoperations	Miyashita T, Kuro M	Department of Anesthesiology, National Cardiovascular Center, Suita, Osaka 565-8565, Japan	Platelets 2001 Jun;12(4):248-53
Comparison of the effects of a cell saver and low-dose aprotinin on blood loss and homologous blood usage in patients undergoing valve surgery	Tempe K, Banerjee A, Virmani S, Mehta N, Panwar S, Tomar A, Ghambeer D, Nigam M	Departments of Anaesthesiology and Cardiothoracic Surgery, G. B. Pant Hospital, New Delhi, India	J Cardiothorac Vasc Anesth 2001 Jun;15(3):326-30
Remaining procoagulant property of wound blood washed by a cell-saving device	Tabuchi N, Sunamori M, Koyama T, Shibamiya A		Ann Thorac Surg 2001 May; 71(5):1749-50
Platelet-rich plasma application in sinus graft surgery: Part I—Background and processing techniques	Lozada JL, Caplanis N, Proussaefs P, Willardsen J, Kammeyer G	Department of Restorative Dentistry, Loma Linda University, Loma Linda, CA	J Oral Implantol 2001; 27(1):38-42
[Removal of lepirudin used as an anticoagulant in mechanical autotransfusion with Cell-Saver 5]	Marx A, von Lupke U, Tessmann R, Lindhoff-Last E	Abteilung für Anästhesie und Intensivmedizin, Berufsgenossenschaftliche Unfallklinik, Frankfurt am Main Germany [Article in German]	Anesthesiol Intensivmed Notfallmed Schmerzther 2001 Mar;36(3):162-6
Fibrinolytic activity and postoperative salvaged untreated blood for autologous transfusion in major orthopaedic surgery	Krohn C, Reikeras O, Bjornsen S, Brosstad F	National Hospital, Centre for Orthopaedics, Oslo, Norway	Eur J Surg 2001 Mar;167(3): 168-72
Autotransfusion decreases blood usage following cardiac surgery – a prospective randomized trial	Dalrymple-Hay J, Dawkins S, Pack L, Deakin C, Sheppard S, Ohri S, Haw M, Livesey S, Monro J	Wessex Cardiothoracic Centre, Southampton, UK	Cardiovasc Surg 2001 Apr; 9(2):184-7
The Role of the Autologous Platelet-Derived Growth Factor in the Management of Decubitus Ulcer	Aminian B, Shams M, Karim-Aghaee B, Soveyd M, Omrani R	Department of Internal Medicine, Shiraz University of Medical Sciences, Shiraz, Iran	[Source unavailable]
Promising Results From a Preliminary Study of Autologous Platelet Gel in Face-lift Surgery	Kent K	1660 Medical Blvd, Suite 100, Naples, FL 34110	Arch Facial Plast Surg 2001 Oct-Dec;3(4):251
Recovery from deep-plane rhytidectomy following unilateral wound treatment with autologous platelet gel: a pilot study	Powell D, Chang E, Farrior E	Division of Facial Plastic and Reconstructive Surgery, Department of Otolaryngology, The Ohio State University, 4100 UHC, 456 W	Arch Facial Plast Surg 2001 Oct-Dec;3(4):245-50
Autologous platelet-rich plasma isolated using the Haemonetics Cell Saver 5 and Haemonetics MCS+ for the preparation of platelet gel	O'Neill E, Zalewski W, Eaton L, Popovsky M, Pivacek L, Ragno G, Valeri C	American Red Cross Blood Services, New England Region, Dedham, MA; USA Naval Blood Research Laboratory, Boston University School of Medicine	Vox Sang 2001 Oct;81(3):172-5

The articles in the following list are provided as additional references for information.

Cobe Cardiovascular is unable to provide copies, but the sources for the articles are given for your convenience in locating full copies.

Title	Authors	Author Affiliation	Source
Intraoperative use of platelet-plasmapheresis in vascular surgery		Dept of Anesthesiology and Pain Medicine, University of California Davis, School of Medicine	J Clin Anesth 2002 Feb;14(1):10-4
Current status of bacterial contamination of autologous blood for transfusion	Sugai Y, Sugai K, Fuse A	Dept of Anesthesia, Ohmihachiman City Hospital, Shiga, Japan	Transfus Apheresis Sci 2001 Jun;24(3):255-9
The use of intraoperative autotransfusion during cranial vault remodeling for craniostyosis	Deva AK, Hopper RA, Landecker A, Flores R, Weiner H, McCarthy JG	Institute of Reconstructive Plastic, NYU Medical Center, New York, NY 10016, USA	Plast Reconstr Surg 2002 Jan;109(1):58-63
Errors during intraoperative cell salvage because of inappropriate wash solutions	Waters JH, Sprung J	Dept of General Anesthesiology, Cleveland Clinic Foundation, Cleveland, OH 44195	Anesth Analg 2001 Dec;93(6):1483-5, table of contents
Activation of plasma coagulation by retransfusion of unwashed drainage blood after hip joint arthroplasty: a prospective study	Duchow J, Ames M, Hess T, Seyfert U	Orthopadische Universitasklinik Homburg-Saar, Germany	J Arthroplasty 2001 Oct;16(7):844-9
Autologous transfusion of shed mediastinal blood after coronary artery bypass grafting and bacterial contamination	Andreason AS, Schmidt H, Jarlov JO, Skov R	Dept of Anesthesiology, Gentofte Hospital, Copenhagen, Denmark	Ann Thorac Surg 2001 Oct;72(4):1327-30
Alveolar ridge and sinus augmentation utilizing platelet-rich plasma in combination with freeze-dried bone allograft: case series	Kassolis JD, Rosen PS, Reynolds MA	Dept of Periodontics, University of Maryland, Dental School, Baltimore, MD, USA	J Periodontol 2000 Oct;71(10):1654-61
Correlation of platelet concentration in platelet-rich plasma to the extraction method, age, sex, and platelet count of the donor	Weibrich G, Kleis WK, Kunz-Kostomanolakis M, Loos AH, Wagner W	Clinic for Oral and Maxillofacial Surgery, Johannes Gutenberg University, Mainz, Germany	Int J Oral Maxillofac Implants 2001 Sep-Oct;16(5):693-9
The use of autologous platelet-rich plasma (platelet gel) and autologous platelet-poor plasma (fibrin glue) in cosmetic surgery	Man D, Plosker H, Winland-Brown JE	Aesthetic Plastic Surgery and Laser Center, Florida Atlantic University, Boca Raton, FL, USA	Plast Reconstr Surg 2001 Jan;107(1):229-37; discussion 238-9